



A case of a giant ectopic parathyroid adenoma diagnosed on 99mTc-MIBI scintigram

Dr. Yousuf Zein-alabideen, Nuclear Medicine - KAASH - TAIF/KSA

Received: 26-03-2018 / Revised Accepted: 30-05-2018 / Published: 02-06-2018

ABSTRACT

We report a case of a 26-year-old man who presented 3 years back with signs and symptoms of severe hypercalcemia and osteoporosis, including muscle spasm and bone deformity and was admitted to hospital several times. Laboratory investigations were consistent with primary hyperparathyroidism (PHPT). 99m Tc-MIBI scintigraphy demonstrated a very big parathyroid adenoma in the mediastinum. Neck and chest computerized tomography scanning confirmed the diagnosis and showed the anatomical details.

Keywords: Hypercalcemia, Tc-MIBI and Osteoporosis

Address for Correspondence: Dr. Yousuf Zein-alabideen MSc, Nuclear Medicine -Radiology Dept. King AbdulAziz Specialist Hospital - TAIF/KSA. email address: yusufzein13@gmail.com

How to Cite this Article: Dr. Yousuf Zein-alabideen. A case of a giant ectopic parathyroid adenoma diagnosed on 99mTc-MIBI scintigram. Int J Res Health Sci 2018; 6(3): 1-5.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, which allows adapt, share and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. 

INTRODUCTION

Primary hyperparathyroidism (PHPT) is a common endocrine disorder characterized by hypercalcemia and excessive secretion of parathyroid hormone (PTH) [1]. PHPT is most commonly caused by a single adenoma of the parathyroid gland. Patients with PHPT tend to develop complications such as reduction of bone mineral density, nephrolithiasis, and gastric ulcer, which may impair quality of life [1, 2]. In the management of PHPT, parathyroidectomy of the abnormal gland is the gold standard for effective treatment. Generally, most parathyroid adenomas remain relatively small, measuring under a few centimeters and weighing less than 1g [3]. Large or giant parathyroid adenomas are seldom seen in patients with PHPT [4], and in such cases differential diagnosis is necessary to rule out malignancy.

Here, we report a case of hyperparathyroidism due to a giant ectopic parathyroid adenoma with complication of chronic hyperparathyroidism and hypercalcemia.

CASE REPORT

A 26-year-old man who had a long history of muscle spasm in hands followed by abnormal gait and was admitted several times to hospital. investigations revealed hypercalcemia. Computed tomography (CT) to skull had shown the presence of multiple osteolytic changes (Fig1). Bone scan and bone densitometry show vertebral scoliosis, generally reduced bone radiotracer uptake and osteoporosis. Nuclear Medicine parathyroid scintigraphy was done using ^{99m}Tc -sestamibi (single isotope method). It showed a very big parathyroid adenoma in the mediastinum (Fig4). Chest CT showed a big elongated retrosternal mass sized 34x 40 mm, positioned in the superior mediastinum below sternum and just to the right of the manubrio-sternal junction (Fig 2 & 3).

DISCUSSION

This is an unusual case of hyperparathyroidism due to a large parathyroid adenoma. This patient presented with signs and symptoms of hypercalcaemia long time back. and was admitted several times to hospital due to complications.

Unfortunately this young man suffered for a quite long time before being diagnosed, and this possibly due to absence of a visible neck swelling on clinical examination. An ectopic parathyroid adenoma should always be considered in such cases especially on the ground of an evident hyperparathyroidism proved by laboratory findings. Parathyroid adenomas usually measure less than 2 cm and weigh less than 1g. Parathyroid carcinomas should always be considered in cases of giant parathyroid adenomas measuring more than 2 cm[7].


Parathyroid cysts or cystic adenomas often show large parathyroid ones. In our case, no signs of malignancy, such as presence of capsular invasion, angioinvasion, and invasion of the surrounding structures, were observed by morphological analysis. In our case, the weight or size of the adenoma are correlated with the functional status of the gland and the severity of biochemical abnormalities, i.e. his large adenoma was associated with a severe form of primary hyperparathyroidism [9]. However, in one reported case of giant adenoma, there was no correlation between size and functional status [10].

As demonstrated in our case, ^{99m}Tc -MIBI-radionuclide imaging has been used in the detection and localization of the parathyroid adenoma, however in the rare occasions of intrathyroid parathyroid adenoma (Incidence: true one is 0.7%, and partial one is 1.9% [12]), radionuclide imaging may miss the pathologic gland [13]. Generally, other different imaging techniques, such as high resolution ultrasonography, CT, arteriography, venous sampling, and magnetic resonance imaging, have been used for detection of the abnormal parathyroid glands [14, 15].


In summary, we report an unusual case of hyperparathyroidism due to a large ectopic parathyroid adenoma positioned in the superior mediastinum.

CONCLUSION


Ectopic parathyroid adenomas can reach a very big size and this could be strongly correlated with the functional status, mostly; as in our case.




(Fig.1)



(fig. 2)



(fig. 3)



(Fig. 4)

REFERENCES

1. Neagoe R. M., Sala D. T., Borda A., Mogoanta C. A., Muhlfay G. Clinicopathologic and therapeutic aspects of giant parathyroid adenomas - three case reports and short review of the literature. *Romanian Journal of Morphology and Embryology*. 2014;55:669–674.
2. Bilezikian J. P., Potts J. T., Jr., El-Hajj Fuleihan G., et al. Summary statement from a workshop on asymptomatic primary hyperparathyroidism: a perspective for the 21st century. *The Journal of Clinical Endocrinology and Metabolism*. 2002;87(12):5353–5361.
3. Bringhurst F. R., Demay M. B., Kronenberg H. M. Hormones and disorders of mineral metabolism. In: Wilson J. D., Foster D. W., Kronenberg H. M., Larsen P. R., editors. *Williams Textbook of Endocrinology*. 9th. Philadelphia, USA: W.B. Saunders Company; 1998. pp. 1155–1209.

4. Krishnamurthy A., Raghunandan G., Ramshankar V. A rare case of giant parathyroid adenoma presenting with recurrent episodes of pancreatitis. *Indian Journal of Nuclear Medicine*. 2016;31(1):36–38.
5. 8. Shah V. N., Bhadada S. K., Bhansali A., et al. Effect of gender, biochemical parameters and parathyroid surgery on gastrointestinal manifestations of symptomatic primary hyperparathyroidism. *Indian Journal of Medical Research*. 2014;139:279–284.
6. Felderbauer P., Karakas E., Fendrich V., Lebert R., Bartsch D. K., Bulut K. Multifactorial genesis of pancreatitis in primary hyperparathyroidism: Evidence for protective (PRSS2) and destructive (CTRC) genetic factors. *Experimental and Clinical Endocrinology and Diabetes*. 2011;119(1):26–29.
7. Araujo Castro M., López A. A., Fragueiro L. M., García N. P. Giant parathyroid adenoma: differential aspects compared to parathyroid carcinoma. *Endocrinology, Diabetes & Metabolism Case Reports*. 2017
8. Ahmad M., Almohaya M., Al Johani N., Almalki M. Intrathyroidal Parathyroid Cyst: An Unusual Neck Mass. *Clinical Medicine Insights: Endocrinology and Diabetes*. 2017;10(0)
9. Zamboni W. A., Folse R. Adenoma weight: A predictor of transient hypocalcemia after parathyroidectomy. *The American Journal of Surgery*. 1986;152(6):611–615.
10. Power C., Kavanagh D., Hill A. D. K., O'Higgins N., McDermott E. Unusual presentation of a giant parathyroid adenoma: Report of a case. *Surgery Today*. 2005;35(3):235–237.
11. O'Doherty M. J., Kettle A. G., Wells P., Collins R. E. C., Coakley A. J. Parathyroid imaging with technetium-99m-sestamibi: Preoperative localization and tissue uptake studies. *Journal of Nuclear Medicine*. 1992;33(3):313–318.
12. Goodman A., Politz D., Lopez J., Norman J. Intrathyroid parathyroid adenoma: Incidence and location - The case against thyroid lobectomy. *Otolaryngology - Head and Neck Surgery*. 2011;144(6):867–871.
13. Bahar G., Feinmesser R., Joshua B.-Z., et al. Hyperfunctioning intrathyroid parathyroid gland: A potential cause of failure in parathyroidectomy. *Surgery*. 2006;139(6):821–826.
14. Krubsack A. J., Wilson S. D., Lawson T. L., et al. Prospective comparison of radionuclide, computed tomographic, sonographic, and magnetic resonance localization of parathyroid tumors. *Surgery*. 1989;106(4):639–644.
15. Bhansali A., Masoodi S. R., Bhadada S., Mittal B. R., Behra A., Singh P. Ultrasonography in detection of single and multiple abnormal parathyroid glands in primary hyperparathyroidism: Comparison with radionuclide scintigraphy and surgery. *Clinical Endocrinology*. 2006;65(3):340–345.